



Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Disease

journal homepage: www.elsevier.com/locate/apjtd

Document heading

doi: 10.1016/S2222-1808(14)60708-8

© 2014 by the Asian Pacific Journal of Tropical Disease. All rights reserved.

Medicinal plants and secondary metabolites for diabetes mellitus control

Mahmoud Bahmani¹, Hannaneh Golshahi², Kouros Saki³, Mahmoud Rafieian-Kopaei^{4*}, Bahram Delfan¹, Tahereh Mohammadi⁵

¹Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran

²Department of Pathology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

³Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁴Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

⁵Tarbiat Modares University, Tehran, Iran

ARTICLE INFO

Article history:

Received 8 Jul 2014

Received in revised form 1 Aug 2014

Accepted 19 Aug 2014

Available online 2 Sep 2014

Keywords:

Diabetes

Medicinal plants

Herbal metabolites

ABSTRACT

Diabetes mellitus is one of the most common and complex problems of modern societies which has caused many economic and social problems. Because diabetes has no definite treatment, the use of traditional medicine seems to be an appropriate solution to control and manage it. Studies revealed that *Vaccinium Arctostaphylos* L., *Securigera securidaca* L., *Gymnema sylvestre* L., *Atriplex halimus* L., *Camellia sinensis* L., *Ginkgo biloba* L., *Mamordica charantia* L., *Citrullus colocynthis* (L.) Schrad., *Allium cepa* L., *Allium sativum* L., *Silybum marianum* (L.), Gaertn and *Trigonella foenum graecum* L. are effective against diabetes. Flavonoids, quercin, metformin, quinalizidine, anthocyanin, catechin and flavone, phenylpropanoids, lipoic acid and coumarin metabolites were introduced major impact on diabetes. With regard to the study of plants and their metabolites and the mechanisms of their influence, it is clear that these plants have the potential to reduce blood sugar and diabetes and be considered as candidates for preparing new drugs. Combination of plants extracts or their components may also have synergistic effects to better act on diabetes.

1. Introduction

Diabetes mellitus is one of the most common and complex problems of modern societies which has caused many economic and social problems. Type I diabetes results from defects in insulin secretion, while the pathogenesis of type II diabetes is associated with a course of progressive insulin resistance in liver and peripheral tissues, reduced β -cell mass and/or impaired insulin secretion[1,2]. Diabetes mellitus causes changes in metabolism of carbohydrates, fats and proteins which result in hyperglycemia, glycosuria, hyperlipidemia and also atherosclerosis[3–5]. Hyperglycemia due to diabetes is associated with long-term damage, destruction and failure of function of various organs[6,7]. One of these organs is the liver and a range of disorders from

fatty change to liver cirrhosis and cancer can occur[8]. Liver is a non-insulin dependent organ and has an important role in lipid and glucose homeostasis[9]. In addition, studies showed tissue damages caused by diabetes can be attributed to oxidative stress[10–13]. Vascular complications of diabetes, including retinopathy, neuropathy, nephropathy and macrovascular disorders are considered as the major causes of mortality in diabetic patients[14–17].

Long-term increase of glucose in diabetes is thought to be the main cause of some disorders like microangiopathy, macroangiopathy, poor antioxidant defense system and impaired lipid metabolism. These disorders can cause short-term and long-term effects which can cause physiological damage to various organs of the body[18,19]. Delayed complications of diabetes such as nephropathy, retinopathy, cardiovascular complications, neuropathy, skin ulcers, high blood pressure and weight gain are more common and more researches have been done about them[20,21]. The number of diabetic patients referred to physicians is increasing significantly. Diabetes affects more than 140 million people in the West countries and the number will reach 300 million by 2025[22].

*Corresponding author: Prof. Mahmoud Rafieian-Kopaei, Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran.

Tel: 0098 913 1811842

Tel/Fax: 0098 381 3349509

E-mail: rafieian@yahoo.com

Foundation Project: Supported by research grant from Deputy for Research and Technology of Lorestan University of Medical Sciences (No. 445/07).

Synthetic drugs are used to treat or prevent diabetes, but they have side effects[23]. Due to this issue and their high costs, nowadays interests in finding effective natural ingredients has increased. In recent decades, many studies have been done on the therapeutic properties of herbs and these elements have been proposed for the treatment of various hard curable diseases such as atherosclerosis[12,24], cardiovascular diseases[25,26], neurological disorders[27,28] and cancer[29,30]. These conditions involve many changes, including alterations in redox state[31,32], and medicinal plants with antioxidant activity have the ability to counteract these situations and always been considered as a healthy source of health promotion[11,33].

Before the discovery of anti-diabetic drugs and insulin, diabetic patients used medicinal plants and traditional medicine[34]. Plants may operate through different mechanisms that effect on blood sugar. Some of them may have insulin kinase[35], and some other may inhibit insulinase activity[36], and others may increase reconstruction of pancreatic β cells[37]. Fibers of plants may also interfere with the absorption of carbohydrates and thus have an effect on blood glucose[38].

There are known herbs and secondary metabolites that are effective in diabetes management and control. In this paper the most effective herbs and plant metabolites for diabetes mellitus will be discussed.

2. Medicinal plants and metabolites on diabetes

2.1. The most effective herbs for diabetes

2.1.1. *Vaccinium arctostaphylos* L.

In traditional medicine this plant is used for treatment of diabetes. This herb contains anthocyanoside myrtillin which has similar mechanism with insulin[39].

2.1.2. *Securigera securidaca* L.

Securigera securidaca L. has significant effect on blood glucose levels in a mouse model of diabetes. It is a native plant of Iran and has a different mechanism from effect of glibenclamide[40].

2.1.3. *Gymnema sylvestre* L. (*G. sylvestre*)

This herb is effective on both types of diabetes by inhibition of absorption of glucose in the digestive tract resulting in glucose excreted and also by reconstruction of damaged beta cells[41,42].

2.1.4. *Atriplex halimus* L.

In animal models it can exerts a significant effect on blood sugar. This herb contains fiber, protein and trace elements such as chromium. Studies showed that administration of three grams of dried leaves of the plant reduces blood sugar in type II diabetics[38].

2.1.5. *Camellia sinensis* L.

Green tea contains flavanols like catechin, epicatechin, epigallocatechin and gallocatechin. These compounds are antioxidants. Powdered green tea increases insulin

resistance. Also, polyphenols in this herb have antioxidant mechanism and have antidiabetic properties[43–45].

2.1.6. *Ginkgo biloba* L.

This herb contains flavoglycosides. It has effect on complications of diabetes and increases the blood flow and indirectly can be effective on diabetic wound healing because of increased blood circulation[46].

2.1.7. *Mamordica charantia* L.

Raw fruit, dried extract, and its juice are effective on diabetes. This herb contains polypeptide which is similar in structure to insulin and reduces blood sugar levels in diabetics Type II[47].

2.1.8. *Citrullus colocynthis* (L.) Schrad.

According to the traditional botanical knowledge as well as clinical studies, *Citrullus colocynthis* fruit has anti-diabetic effects[48].

2.1.9. *Allium cepa* L.

Allyl propyl disulfide is a sulfur compound in onions that have glucose lowering effects[49].

2.1.10. *Allium sativum* L.

Allyl disulfide oxide as the active ingredient in garlic has a blood sugar lowering effects[38].

2.1.11. *Silybum marianum* (L.) Gaertn

Silybin and silydianin which are flavonoids in this herb, can increase cellular uptake of glucose[50].

2.1.12. *Trigonella foenum graecum* L.

The seeds of this herb contain bioactive substances such as trigonelline, coumarin and nicotinic acid. Seeds have antidiabetic properties. Fifteen grams of soaked powdered seeds can reduce blood glucose in type II diabetic patients. Amino acids and fiber, have been introduced as antidiabetic agents[51].

2.2. The most common metabolites influencing diabetes

Although there is a wide list of natural anti-diabetes drugs, however, citing all natural antidiabetic plants and their components is out of the scope of this paper. We tried to summarize the most biologically active secondary metabolites with anti-diabetes activity in Table 1 discussing some of them below[52].

2.2.1. Flavonoids

Flavonoids are constituents of fruits, vegetables and herbs. Flavonoids (quercetin) can stimulate insulin secretion. Quercetin inhibits aldolase reductase in diabetic patients. Quercetin exists in cabbage, green tea, apple, onion, beans and nuts, and such herbs like *Ginkgo biloba*[53].

Metformin or dimethyl-biguanide is an oral anti-diabetic herbal ingredient which is derived from plant *Galega officinalis*. It has a wide variety of other therapeutic effects[21,54].

Table 1

The most biologically active secondary metabolites with anti-diabetes activity.

Natural compounds class	Name	Mechanism of action and uses
Alkaloids	Casuarine 6-O- α -glucoside	α -Glucosidase inhibitor
	Isoquinoline alkaloids: schulzeines A, B and C	α -Glucosidase inhibitor
	Tecomine, 5 β -hydroxyskitanthine, boschniakine	Stimulate basal glucose uptake rate in rat adipocytes
	Two new pyrrolidine alkaloids : radicamines A and B	α -Glucosidase inhibitor
	Three quinolizidine alkaloids: javaberine A, javaberine A hexaacetate and javaberine B hexaacetate	Inhibitors of TNF- α production by macrophages and fat cells.
Flavonoids	Three quinolizidine alkaloids: lupanine, 13- α -hydroxylupanine, 17-oxo-lupanine	Dietary supplement for prevention of diabetes Glucose-induced insulin release enhancement from isolated rat islet cells which was dependent on the glucose concentration
	Five isoquinoline: alkaloids berberine, chloride berberine, sulfate berberine, iodide, palmatine sulfate, palmatine chloride	Aldose reductase inhibitor
	6-Hydroxyapigenin, 6-hydroxyapigenin-7-O- β -D-glucopyranoside, 6-hydroxyluteolin- 7-O- β -D-glucopyranoside, 6-hydroxyapigenin-7-O- (6-O-feruloyl)- β -D-glucopyranoside, 6-hydroxyluteolin-7-O-(6-O-feruloyl)- β -D-glucopyranoside	α -Glucosidase inhibitor
	Myricitrin I, II, III, IV and V	Aldose reductase inhibitory activity.
	Quercetin 3-O- α -L-arabinopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside, Kaempferol 3-O- β -D-glucopyranoside (astragalin), quercetin 3-O- β -D-glucopyranoside (isoquercitrin)	Glycation inhibitors
Terpenes	Lactucain A, B and C	Moderate lowering of plasma glucose
	3,22-Dihydroxyolean-12-en-29-oic acid, tingenone, tingenine B, regeol A, triptocalline A, mangiferin	Aldose reductase inhibitory activity
	Centellasaponin A	Aldose reductase inhibitory activity
	Abietane-type diterpenoids: danshenols A and B, dihydrotanshinone I, tanshinone I, cryptotanshinone, tanshinone IIA (-)-danshexinkun A	Aldose reductase inhibitory activity
	Triterpenedehydrotrametenolic acid	Insulin sensitizer
Phenolics	Corosolic acid (GlucosolTM)	Glucose transport-stimulating activity
	7'-(3',4'-Dihydroxyphenyl)-N-[(4 methoxyphenyl)ethyl] propenamide, 7'-(4'-hydroxy-3'methoxyphenyl)-N-[(4butylphenyl)ethyl] propenamide	α -Glucosidase inhibitor
	(7S,8S)-syringoylglycerol 9-O- β -D-glucopyranoside (7S,8S)-Syringoylglycerol-9-O-(6'-O-cinnamoyl)- β -D-glucopyranoside	α -Glucosidase inhibitor
	Tetra- and penta-O-galloyl- β -D-glucose	Potent aldose reductase inhibitory activities.

2.2.2. Quinolizidine alkaloids

Some alkaloids, such as sparteine and lupanine are derived from quinolizidine which is a heterocyclic compound and contains nitrogen. About 150 alkaloids of quinolizidine group are known in various species of *Lupinus*. These compounds can stimulate insulin secretion in a glucose-dependent manner and partly through inhibition of adenosine triphosphate dependent potassium channels in the beta cells of the pancreas[55].

Metformin reduces blood glucose and may eliminate insulin sensitivity, but in the absence of insulin (as in type I diabetes) it is ineffective. Mechanism of metformin in non-insulin dependent diabetic patients is attributed to reducing the output of hepatic glucose and increasing peripheral glucose uptake[56].

Flavonoids are found in many plants: anthocyanins (petunidin in raspberry, peonidin in strawberry, pelargonidin in tea, delphinidin in red grapes, cyanidin in cherry and malvidin in red wine), catechin (epicatechin in red wine and catechin in tea), flavone (quercetin in potato skins, apigenin in olive, luteolin in parsley, sabelin the celery, myricetin in broccoli and rutin in lettuce), flavanone (taxifolin, naringenin, naringin, fisetin and hesperetin in peel and fruit of citrus)[57].

Phenylpropanoids, a class of organic compounds derived from phenylalanine are a diverse group of plant secondary metabolites. They include flavonoids, stilbenoids and lignins. Different compounds in this group have hypoglycaemic properties[58].

Lipoic acid exists in green vegetables such as broccoli,

spinach and animal products of ruminants such as the heart, liver, kidney and skeletal muscles. Clinical studies have shown that the lipoic acid improves glucose metabolism in type II diabetic patients by stimulating the uptake of glucose by the rapid transfer through glucose channels (glucose transporter type 4) from intracytosol vesicles to plasma membrane[59–61].

Coumarin is the major compound of turmeric which is also effective on diabetes mellitus induced by streptozotocin and nicotine amide. The mechanism is via increased hexokinase activity and glucose-6-phosphate dehydrogenase, namely increase in glucose decomposition and the significant reduction in titer of enzymes glucose 6-phosphatase and fructose-1, 6-bisphosphatase (decrease gluconeogenesis) [62].

3. Conclusions

Nowadays, the data on medicinal plants and their biological activities are tremendously increasing[34,63,64]. However, it is impractical to specify the performance of a mixture which present in the plant extracts comprising a wide range of phytochemical constituents, to only a single compound from that extract[65]. Metabolites might act as lead compounds for the discovery of different new classes of possibly potent and safe anti-diabetic agents. However, attention should be given for the identification of the typical modes of action of their extracts and the isolated pure compounds[66].

Many classes of secondary metabolites of herbal medicines, such as alkaloids, phenolics, flavonoids, terpenoids and others have been shown promising anti-diabetic activities. These constituents may be considered as a promising source of hypoglycemic agents with minimal side effects.

Finding the promising candidates for the treatment of diabetes mellitus could be achieved through evaluation of the anti-diabetic properties of different medicinal plant extracts, their fractions and isolated components, followed by preliminary phytochemical screening, and their toxicities^[23,67]. Although various compounds with different mechanism of actions have been shown to reduce blood glucose, however, antioxidant activity of these plants has a crucial role in their anti-diabetes actions^[10,11,13,32,33,68].

Oxidative stress has been implicated not only in diabetes mellitus, but also in a wide variety of pathological conditions such as cancer^[69,70], neurological disorders^[71,72], ischemia/reperfusion^[73,74]. These conditions involve many changes, including impaired glucose tolerance, alterations in the thiol/disulphide redox state, and activation of inflammatory processes^[11,75,76]. They are also associated with increased activity of nicotinamide adenine dinucleotide phosphate oxidase or induction of xanthine oxidase with subsequent increase in oxidative stress during chronic inflammation and the generation of damage, particularly during diabetes and ischemia/reperfusion. Exposure to agents that increase the oxidative stress might accelerate the initiation of pathological changes such as diabetes. Therefore, antioxidant activity evaluation of the plants will give a preliminary prediction for their anti-diabetic activities^[77,78].

Pharmacological evaluation to recognize the most active candidates will be determined through measuring different biochemical parameters such as serum glucose, insulin, glycosylated hemoglobin, lipid profile, serum urea and creatinine, plasma aspartate transaminase and alanine transaminase. The anti-diabetic properties will also be further confirmed through microscopical examinations of the pancreatic sections^[4,16,18].

Botany traditional knowledge and clinical knowledge are linked together. A lot of clinical trials have taken knowledge of botany source. Many of the traditional knowledge of medicinal plants have been routinely taken in the treatment of various diseases from this knowledge^[79–87]. Therefore, the use of more traditional resources have been recently recommended until further herbs in the treatment of diabetes presented, and with clinical research production of new herbal drugs were developed.

Medicinal plants are still considered as a promising source for drug discovery and are playing a pivotal role in drug development programs. Moreover, many medicinal plants provide a rich mine for bioactive chemicals that are mostly free from undesirable adverse effects and of powerful pharmacological actions.

Much effort should be afforded to optimize a procedure for antidiabetic screening of different plants' extracts as well as isolating bioactive compounds for the discovery of new natural herbal antidiabetic drugs that can be used as alternatives to synthetic oral hypoglycemic drugs with less or even no prominent side effects.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

This study was conducted in cooperation with the Deputy for Research and Technology of Lorestan University of Medical Sciences. This research was supported by research grant (No. 445/07) from Deputy for Research and Technology of Lorestan University of Medical Sciences.

References

- [1] Srinivasan K, Ramarao P. Animal models in type 2 diabetes research: an overview. *Indian J Med Res* 2007; **125**: 451–472.
- [2] Matthaie S, Stumvoll M, Kellerer M, Häring HU. Pathophysiology and pharmacological treatment of insulin resistance. *Endocr Rev* 2000; **2**: 585–618.
- [3] Behradmanesh S, Horestani MK, Baradaran A, Nasri H. Association of serum uric acid with proteinuria in type 2 diabetic patients. *J Res Med Sci* 2013; **18**: 44–46.
- [4] Baradaran A, Madihi Y, Merrikhi A, Rafieian-Kopaei M, Nasri H. Serum lipoprotein (a) in diabetic patients with various renal function not yet on dialysis. *Pak J Med Sci* 2013; **29**: 354–357.
- [5] Mirhoseini M, Baradaran A, Rafieian-Kopaei M. Medicinal plants, diabetes mellitus and urgent needs. *J HerbMed Pharmacol* 2013; **2**(2): 53–54.
- [6] Lyra R, Oliveira M, Lins D, Cavalcanti N. [Prevention of type 2 diabetes mellitus]. *Arq Bras Endocrinol Metabol* 2006; **50**(2): 239–249. Portuguese.
- [7] Baradaran A, Madihi Y, Merrikhi A, Rafieian-Kopaei M, Nasri H. Serum lipoprotein (a) in diabetic patients with various renal function not yet on dialysis. *Pak J Med Sci* 2013; **29**(1): 354–357.
- [8] Nasri H, Sahinfard N, Rafieian M, Rafieian S, Shirzad M, Rafieian-Kopaei M. Effects of *Allium sativum* on liver enzymes and atherosclerotic risk factors. *J HerbMed Pharmacol* 2013; **2**(2): 23–28.
- [9] Moller DE. New drug targets for type 2 diabetes and the metabolic syndrome. *Nature* 2001; **414**(6865): 821–827.
- [10] Nasri H, Rafieian-Kopaei M. Tubular kidney protection by antioxidants. *Iran J Public Health* 2013; **42**(10): 1194–1196.
- [11] Nasri H, Rafieian-Kopaei M. Protective effects of herbal antioxidants on diabetic kidney disease. *J Res Med Sci* 2014; **19**(1): 82–83.
- [12] Madihi Y, Merrikhi A, Baradaran A, Rafieian-Kopaei M, Shahinfard N, Ansari R, et al. Impact of sumac on postprandial high-fat oxidative stress. *Pak J Med Sci* 2013; **29**(1): 340–345.
- [13] Rafieian-Kopaei M, Beyki A, Nasri H. [Antioxidant therapy in hemodialysis patients]. *J Isfahan Med School* 2013; **31**(261): 1897–1900. Persian.
- [14] Rafieian-Kopaei M, Nasri H. Comment on: Is the renoprotective effect of erythropoietin in chronic kidney disease a myth? *J Formos Med Assoc* 2014; **113**(1): 62.
- [15] Rafieian-Kopaei M, Ghaed-Amini F, Nasri H. Erythropoietin has kidney protective efficacy beyond stimulating erythropoiesis. *J Isfahan Med School* 2013; **31**: 619–622.
- [16] Nasri H, Rafieian-Kopaei M. Preventive and curative effect of

- garlic on nephrotoxic effect of gentamicin in rat. *J Babol Univ Med Sci* 2014; **16**(2): 42–48.
- [17] Rafieian-Kopaei M, Ghobadi Sh, Nasri H. [Hyperuricemia aggravates IgA nephropathy]. *J Isfahan Med School* 2013; **31**(253): 15518–1521. Persian.
- [18] Nasri H, Rafieian-Kopaei M. Medicinal plants and new concerns in statin consumption. *Iran J Public Health* 2013; **42**(9): 1071–1072.
- [19] Rafieian-Kopaei M, Baradaran A, Nasri H. Significance of extracapillary proliferation in IgA-nephropathy patients with regard to clinical and histopathological variable. *Hippokratia* 2013; **17**(3): 258–261.
- [20] Amini GF, Rafieian-Kopaei M, Nematbakhsh M, Baradaran A, Nasri H. Ameliorative effects of metformin on renal histologic and biochemical alterations of gentamicin-induced renal toxicity in Wistar rats. *J Res Med Sci* 2012; **17**(7): 621–625.
- [21] Nasri H, Baradaran A, Ardalan MR, Mardani S, Momeni A, Rafieian-Kopaei M. Bright renoprotective properties of metformin: beyond blood glucose regulatory effects. *Iran J Kidney Dis* 2013; **7**(6): 423–428.
- [22] King H, Aubert RE, Herman WH. Global burden of diabetes, 1995–2025: Prevalence, numerical estimates, and projections. *Diabetes Care* 1998; **21**: 1414–1431.
- [23] Nasri H, Shirzad H. Toxicity and safety of medicinal plants. *J HerbMed Pharmacol* 2013; **2**(2): 21–22.
- [24] Setorki M, Rafieian-Kopaei M, Merikhi A, Heidarian E, Shahinfard N, Ansari R, et al. Suppressing impact of *Anethum graveolens* consumption on biochemical risk factors of atherosclerosis in hypercholesterolemic rabbits. *Int J Prev Med* 2013; **4**(8): 889–895.
- [25] Khosravi-Boroujeni H, Mohammadifard N, Sarrafzadegan N, Sajjadi F, Maghroun M, Khosravi A, et al. Potato consumption and cardiovascular disease risk factors among Iranian population. *Int J Food Sci Nutr* 2012; **63**(8): 913–920.
- [26] Khosravi-Boroujeni H, Sarrafzadegan N, Mohammadifard N, Sajjadi F, Maghroun M, Asgari S, et al. White rice consumption and CVD risk factors among Iranian population. *J Health Popul Nutr* 2013; **31**(2): 252–261.
- [27] Akhlaghi M, Shabani G, Rafieian-Kopaei M, Parvin N, Saadat M, Akhlaghi M. *Citrus aurantium* blossom and preoperative anxiety. *Rev Bras Anesthesiol* 2011; **61**(6): 702–712.
- [28] Roohafza H, Sarrafzadegan N, Sadeghi M, Rafieian-Kopaei M, Sajjadi F, Khosravi-Boroujeni H. The association between stress levels and food consumption among Iranian population. *Arch Iran Med* 2013; **16**(3): 145–148.
- [29] Azadmehr A, Hajiaghae R, Afshari A, Amirghofran Z, Rafieian-Kopaei M, Darani HY, et al. Evaluation of *in vivo* immune response activity and *in vitro* anti-cancer effect by *Scrophularia megalantha*. *J Med Plants Res* 2011; **5**(11): 2365–2368.
- [30] Shirzad H, Shahrani M, Rafieian-Kopaei M. Comparison of morphine and tramadol effects on phagocytic activity of mice peritoneal phagocytes *in vivo*. *Int Immunopharmacol* 2009; **9**(7–8): 968–970.
- [31] Rafieian-Kopaei M, Baradaran A, Rafieian M. Plants antioxidants: from laboratory to clinic. *J Nephropathol* 2013; **2**(2): 152–153.
- [32] Kafash-Farkhad N, Asadi-Samani M, Rafieian-Kopaei M. A review on phytochemistry and pharmacological effects of *Prangos ferulacea* (L.) Lindl. *Life Sci J* 2013; **10**(8s): 360–367.
- [33] Baradaran A, Nasri H, Nematbakhsh M, Rafieian-Kopaei M. Antioxidant activity and preventive effect of aqueous leaf extract of *Aloe vera* on gentamicin-induced nephrotoxicity in male Wistar rats. *Clin Ter* 2014; **165**(1): 7–11.
- [34] Rafieian-Kopaei M, Sewell RDE. The history and ups and downs of herbal medicine usage. *J HerbMed Pharmacol* 2014; **3**(1): 1–3.
- [35] Bailey CJ, Day C. Traditional plant medicine as treatment for diabetes. *Diabetes Care* 1989; **12**: 553–564.
- [36] Abdel-Moneim A, El-Feki M, Salah E. Effect of *Nigella sativa*, fish oil and gliclazide on alloxan diabetic rats 1–biochemical and histopathological studies. *J Egy Ger Soc Zool* 1999; **23**: 237–265.
- [37] Chakravarthy BK, Saroj G, Gambhir SS, Gode KD. Pancreatic beta cell regeneration—a novel antidiabetic mechanism of *Pterocarpus marsupium* roxb. *Indian J Pharmacol* 1980; **12**: 123–128.
- [38] Bever BO, Zahnd GR. Plants with oral hypoglycemic action. *Pharm Biol* 1979; **17**: 139–196.
- [39] Murray M. Bilberry (*Vaccinium myrtillus*). *Am J Nat Med* 1997; **4**: 18–22.
- [40] Hisseinzadeh H, Ramezani M, Danaei AR. Antihyperglycaemic effect and acute toxicity of *Securigera secuidaca* L. seed extracts in mice. *Phytother Res* 2002; **16**: 745–747.
- [41] Shanmugasundaram ER, Rajeswari G, Baskaran K, Rajesh Kumar BR, Radha Shanmugasundaram K, Kizar Ahmath B. Use of *Gymnema sylvestre* leaf extract in the control of blood glucose in insulin dependent diabetes mellitus. *J Ethnopharmacol* 1990; **30**: 281–294.
- [42] Baskaran K, Kizar Ahmath B, Radha Shanmugasundaram K, Shanmugasundaram ER. Antidiabetic effect of a leaf extract from *Gymnema sylvestre* in non-insulin dependent diabetes mellitus patients. *J Ethnopharmacol* 1990; **30**: 295–300.
- [43] Guo Q, Zhao B, Shen S, Hou J, Hu J, Xin W. ESR study on the structure-antioxidant activity relationship of tea catechins and their epimers. *Biochim Biophys Acta* 1999; **1427**: 13–23.
- [44] Asadi Y, Karimi M, Parsaei P, Ezzati S, Khadivi R, Zamiri A, et al. Effects of *Camellia sinensis* ethanolic extract on histometric and histopathological healing process of burn wound in rat. *Middle-East J Sci Res* 2013; **13**(1): 14–19.
- [45] Parsaei P, Karimi M, Asadi SY, Rafieian-Kopaei M. Bioactive components and preventive effect of green tea (*Camellia sinensis*) extract on postlaparotomy intra-abdominal adhesion in rats. *Int J Surg* 2013; **11**(9): 811–815.
- [46] Bauer U. 6-Month double-blind randomized clinical trial of *Ginkgo biloba* extract versus placebo in two parallel groups in patients suffering from peripheral arterial insufficiency. *Arzneimittelforschung* 1984; **34**: 716–720.
- [47] Welihinda J, Arvidson G, Gylfe E, Hellman B, Karlsson E. The insulin-releasing activity of the tropical plant *Momordica charantia*. *Acta Biol Med Ger* 1982; **41**: 1229–1240.
- [48] Rahman MA, Mossa JS, Al-Said MS, Al-Yahya MA. Medicinal plant diversity in the flora of Saudi Arabia 1: a report on seven plant families. *Fitoterapia* 2004; **75**: 149–161.
- [49] Sheela CG, Augusti KT. Antidiabetic effects of S-allyl cysteine sulfoxide isolated from garlic (*Allium sativum* L.). *Indian J Exp Biol* 1992; **30**: 523–526.
- [50] Lirussi F, Beccarello A, Zanette G, De Monte A, Donadon V, Velussi M, et al. Silybin–betacyclodextrin in the treatment of patients with diabetes mellitus and alcoholic liver disease. Efficacy study of a new preparation of an anti-oxidant agent. *Diabetes Nutr Metab* 2002; **15**: 222–231.
- [51] Ribes G, Sauvaire Y, Baccou JC, Valette G, Chenon D, Trimble ER, et al. Effects of fenugreek seeds on endocrine pancreatic secretions in dogs. *Ann Nutr Metab* 1984; **28**: 37–43.
- [52] Singab AN, Youssef FS, Ashour ML. Medicinal plants with potential antidiabetic activity and their assessment. *Med Aromat Plants* 2014; **3**(1): 151.
- [53] Arabbi PR, Genovese MI, Lajolo FM. Flavonoids in vegetable

- foods commonly consumed in Brazil and estimated ingestion by the Brazilian population. *J Agric Food Chem* 2004; **52**: 1124–1131.
- [54] Rafieian-Kopaei M, Baradaran A, Merrikhi A, Nematbakhsh M, Madihi Y, Nasri H. Efficacy of co-administration of garlic extract and metformin for prevention of gentamicin–renal toxicity in wistar rats: A biochemical study. *Int J Prev Med* 2013; **4**(3): 258–264.
- [55] Wink M. Plant breeding: importance of plant secondary metabolites for protection against pathogens and herbivores. *Theor Appl Genet* 1988; **75**: 225–233.
- [56] Bailey CJ. Metformin—an update. *Gen Pharmacol* 1993; **24**: 1299–1309.
- [57] Rice–Evans CA, Miller NJ, Paganga G. Structure–antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic Biol Med* 1996; **20**: 933–956.
- [58] Vogt T. Phenylpropanoid biosynthesis. *Mol Plant* 2010; **3**(1): 2–20.
- [59] Konrad T, Vicini P, Kusterer K, Höflich A, Assadkhani A, Böhles HJ, et al. alpha–Lipoic acid treatment decreases serum lactate and pyruvate concentrations and improves glucose effectiveness in lean and obese patients with type 2 diabetes. *Diabetes Care* 1999; **22**: 280–287.
- [60] Jacob S, Ruus P, Hermann R, Tritschler HJ, Maerker E, Renn W, et al. Oral administration of RAC–alpha–lipoic acid modulates insulin sensitivity in patients with type–2 diabetes mellitus: a placebo–controlled pilot trial. *Free Radic Biol Med* 1999; **27**: 309–314.
- [61] Estrada DE, Ewart HS, Tsakiridis T, Volchuk A, Ramlal T, Tritschler H, et al. Stimulation of glucose uptake by the natural coenzyme alphas lipoic acid/thioctic acid: participation of elements of the insulin signaling pathway. *Diabetes* 1996; **45**: 1798–1804.
- [62] Pari L, Rajarajeswari N. Efficacy of coumarin on hepatic key enzymes of glucose metabolism in chemical induced type 2 diabetic rats. *Chem Biol Interact* 2009; **181**: 292–296.
- [63] Nasri H, Tavakoli M, Ahmadi A, Baradaran A, Nematbakhsh M, Rafieian-Kopaei M. Ameliorative effect of melatonin against contrast media induced renal tubular cell injury. *Pak J Med Sci* 2014; **30**(2): 261–265.
- [64] Asadi SY, Parsaei P, Karimi M, Ezzati S, Zamiri A, Mohammadzadeh F, et al. Effect of green tea (*Camellia sinensis*) extract on healing process of surgical wounds in rat. *Int J Surg* 2013; **11**(4): 332–337.
- [65] Wink M. Evolutionary advantage and molecular modes of action of multi–component mixtures used in phytomedicine. *Curr Drug Metab* 2008; **9**: 996–1009.
- [66] Rafieian-Kopaei M. Medicinal plants and the human needs. *J HerbMed Pharmacol* 2012; **1**(1): 1–2.
- [67] Bnouham M, Ziyat A, Mekhfi H, Tahri A, Legssyer A. Medicinal plants with potential antidiabetic activity—A review of ten years of herbal medicine research (1990–2000). *Int J Diabetes Metab* 2006; **14**: 1–25.
- [68] Sharafati-Chaleshtori R, Sharafati-Chaleshtori F, Rafieian M. Biological characterization of Iranian walnut (*Juglans regia*) leaves. *Turk J Biol* 2011; doi: 10.3906/biy-1005-1.
- [69] Shirzad H, Taji F, Rafieian-Kopaei M. Correlation between antioxidant activity of garlic extracts and WEHI–164 fibrosarcoma tumor growth in BALB/c mice. *J Med Food* 2011; **14**(9): 969–974.
- [70] Shirzad M, Kordyazdi R, Shahinfard N, Nikokar M. Does royal jelly affect tumor cells? *J HerbMed Pharmacol* 2013; **2**(2): 45–48.
- [71] Shirani M, Alibabaei Z, Kheiri S, Shirzad H, Taji F, Asgari A, Rafieian M. Effect of *Euphorbia helioscopia* extract on acute and chronic pain in mice. *J Babol Univ Med Sci* 2011; **13**(4): 14–18.
- [72] Mirzaei MGR, Azimian M, Moezzi M, Vameghi R, Rafieian-Kopaei M. Effect of lamotrigine on prophylaxis of pediatric classic migraine. *Iran J Child Neurol* 2009; **35**: 35–38.
- [73] Rabiei Z, Rafieian-Kopaei M, Heidarian E, Saghaei E, Mokhtari S. Effects of *Zizyphus jujube* extract on memory and learning impairment induced by bilateral electric lesions of the nucleus basalis of meynert in rat. *Neurochem Res* 2013; **39**(2): 353–360.
- [74] Rabiei Z, Rafieian-Kopaei M, Mokhtari S, Alibabaei Z, Shahrani M. The effect of pretreatment with different doses of *Lavandula officinalis* ethanolic extract on memory, learning and nociception. *Biomed Aging Pathol* 2013; **4**: 71–76.
- [75] Asgari S, Keshvari M, Sahebkar A, Hashemi M, Rafieian-Kopaei M. Clinical investigation of the acute effects of pomegranate juice on blood pressure and endothelial function in hypertensive individuals. *ARYA Atheroscler* 2013; **9**(6): 326–331.
- [76] Asgari S, Sahebkar A, Afshani M, Keshvari M, Haghighjooyjavanmard Sh, Rafieian-Kopaei M. Clinical evaluation of blood pressure lowering, endothelial function improving, hypolipidemic and anti-inflammatory effects of pomegranate juice in hypertensive subjects. *Phytother Res* 2013; **28**(2): 193–199.
- [77] Nasri H, Rafieian-Kopaei M. Oxidative stress and aging prevention. *Int J Prev Med* 2013; **4**(9): 1101–1102.
- [78] Rafieian-Kopaei M, Baradaran A, Rafieian M. Oxidative stress and the paradoxical effects of antioxidants. *J Res Med Sci* 2013; **18**(7): 629.
- [79] Bahmani M, Eftekhari Z. An ethnoveterinary study of medicinal plants in treatment of diseases and syndromes of herd dog in southern regions of Ilam province, Iran. *Comp Clin Pathol* 2012; **22**: 403–407.
- [80] Ghasemi Pirbalouti A, Momeni M, Bahmani M. Ethnobotanical study of medicinal plants used by Kurd tribe in Dehloran and Abadan districts, Ilam province, Iran. *Afr J Tradit Complement Altern Med* 2013; **10**(2): 368–385.
- [81] Bahmani M, Rafieian-Kopaei M. Medicinal plants and secondary metabolites for leech control. *Asian Pac J Trop Dis* 2014; **4**(4): 315–316.
- [82] Bahmani M, Farkhondeh T, Sadighara P. The anti-parasitic effects of *Nicotina tabacum* on leeches. *Comp Clin Pathol* 2012; **21**(3): 357–359.
- [83] Bahmani M, Karamati SA, Banihabib E, Saki K. Comparison of effect of nicotine and levamisole and ivermectin on mortality of leech. *Asian Pac J Trop Dis* 2014; **4**(Suppl 1): S477–S480.
- [84] Amirmohammadi M, Khajoenia SH, Bahmani M, Rafieian-Kopaei M, Eftekhari Z, Qorbani M. *In vivo* evaluation of antiparasitic effects of *Artemisia abrotanum* and *Salvia officinalis* extracts on *Syphacia obvelata*, *Aspiculuris tetrapetra* and *Hymenolepis nana* parasites. *Asian Pac J Trop Dis* 2014; **4**(Suppl 1): S250–S254.
- [85] Bahmani M, Abbasi J, Mohsenzadegan A, Sadeghian S, Ahangaran MG. *Allium sativum* L.: the anti-immature leech (*Limnatis nilotica*) activity compared to niclosamide. *Comp Clin Pathol* 2013; **22**: 165–168.
- [86] Bahmani M, Banihabib E, Rafieian-Kopaei M, Gholami-Ahangaran M. Comparison of disinfection activities of nicotine with copper sulphate in water containing *Limnatis nilotica*. *Kafkas Univ Vet Fak Derg*. Forthcoming 2014.
- [87] Bahmani M, Saki K, Rafieian-Kopaei M, Karamati SA, Eftekhari Z, Jelodari M. The most common herbal medicines affecting Sarcomastigophora branches: a review study. *Asian Pac J Trop Biomed* 2014; **4**(12): 930–937.